

Pulmonary Torsion as an Atypical Complication of Congenital Esophageal Atresia Repair—A Case Report and Review of Literature

Maria Koziarkiewicz¹ Anna Taczalska¹ Anna Piaseczna-Piotrowska¹

¹ Department of Pediatric Surgery and Urology, Polish Mother's Health Center Research Institute, Lodz, Poland

Address for correspondence Maria Koziarkiewicz, MD, PhD, Department of Pediatric Surgery and Urology, Polish Mother's Health Research Institute, 93-338 Lodz, Rzgowska 281/289, Poland (e-mail: amaltea79@interia.pl).

Eur J Pediatr Surg Rep 2014;2:43–45.

Abstract

Keywords

- pulmonary torsion
- esophageal atresia
- postoperative complication

Pulmonary torsion is a severe, life-threatening event, rarely occurring in children. We present a case of atypical postoperative complication of esophageal atresia repair in the form of pulmonary torsion comprising the middle lobe of the right lung. Clinical deterioration in the face of normal arterial blood gases should rise a high index of suspicion for pulmonary torsion. Early diagnosis is crucial, treatment is surgical, detorsion if possible, or if the lobe is not viable—resection.

Introduction

Esophageal atresia is an anatomical and physiological disorder of the esophagus requiring urgent surgical intervention. The most common early complications of esophageal atresia repair include leakage of the anastomosis, recurrent tracheoesophageal fistula, and anastomotic stricture. We present a case of atypical postoperative complication in the form of pulmonary torsion comprising the middle lobe of the right lung.

Case Report

A 3,260 g preterm female neonate, delivered at 38 weeks' gestation, evaluated at 7/8 points in Apgar scale, was referred to our department with prenatally diagnosed esophageal atresia. On contrast, chest X-ray performed directly after birth the presence of congenital defect—esophageal atresia with tracheoesophageal fistula—was confirmed (the upper end of esophagus visible at the level of Th4–5, dilated intestinal loops filled with air). The child was operated on the second day of life by the right laterodorsal thoracotomy. The tracheoesophageal fistula was dissected and sutured, and esophago-esophageal anastomosis was performed. Before closure, the lungs were aerated by forced ventilation, continued until all collapsed regions fully expanded. During post-

operative course, on control chest X-rays, there was atelectasis in the upper lobe and the middle lobe of the right lung (►Fig. 1). On ultrasound examination, no pleural effusion was noted. On the sixth postoperative day, contrast-enhanced computed tomographic (CT) scan was performed and revealed abrupt termination of the bronchial airways with atypical course of horizontal interlobar fissure (►Fig. 2). Right pulmonary artery was visible, its lobar branches hardly visible; right main bronchus showed typical diameter for approximately 7 mm from the bifurcation, then narrowed to approximately 1.5 mm (compression from outside?) and continued as a single bronchus to the lower lobe. There were neither any branches to the upper lobe nor to the middle lobe. Thus, bronchography was performed, which showed obstruction of the upper lobe bronchus and bronchus intermedius and the middle lobe bronchus of the right lung (►Fig. 3).

The patient was qualified for tracheobronchoscopy, which confirmed occlusion of the upper and middle lobar bronchi (trachea—wide, smooth, site of sutured fistula wide, patent, tracheal bifurcation normal. Main bronchus: left, normal; right, narrowing at ~ 10 mm from bifurcation, with a small fissure in the sagittal plane). The neonate was qualified for second thoracotomy on the ninth day of life. On exploration, massive infarction of the upper and middle lobe of the right lung was revealed. Anastomosis site and site of

received
May 31, 2013
accepted after revision
October 1, 2013
published online
January 16, 2014

DOI <http://dx.doi.org/10.1055/s-0033-1361834>.
ISSN 2194-7619.

© 2014 Georg Thieme Verlag KG
Stuttgart · New York

License terms



response and multiorgan failure.^{1,13,14} In the two described cases of detorsion, one patient recovered completely,¹ whereas in the second case the lung remained shrunken and nonfunctioning.⁶ In this case, detorsion allowed for better assessment of the lobe viability, and decrease in edema and hemostasis were observed. In the case of our patient, the torsion was probably because of an anatomic anomaly—absence of parenchymal bridges between contiguous lobes. Earlier reoperation would have probably been beneficial, but as this complication is extremely rare in the pediatric population, diagnostic process may take longer. Persistent atelectasis of a lung/lobe after thoracotomy should call attention to a possibility of torsion.

Conclusions

The following can be concluded from this case report:

1. Pulmonary torsion is a severe, life-threatening event, rarely occurring in children.
2. Early diagnosis (X-ray/CT/bronchography/bronchoscopy) may be lifesaving. Clinical deterioration in the face of normal arterial blood gases should raise a high index of suspicion for diagnosis.
3. Treatment is surgical, detorsion if possible, or if the lobe is not viable—resection.

References

- 1 Oliveira C, Zamakhshary M, Abdallah MR, et al. Lung torsion after tracheoesophageal fistula repair: a case report and review of literature. *J Pediatr Surg* 2007;42(11):E5–E9
- 2 Alberti D, Borsellino A, Migliazza L, et al. Pulmonary torsion after cardiac surgery in two infants: review of pediatric literature. *J Pediatr Surg* 2004;39(11):1719–1723
- 3 Stratemeier EH, Barry JW. Torsion of the lung following thoracic trauma; a case report. *Radiology* 1954;62(5):726–727
- 4 Parks RE. Traumatic torsion of the lung. *Radiology* 1956;67(4):582–583
- 5 Selmonosky CA, Flege JB Jr, Ehrenhaft JL. Torsion of a lobe of the lung due to blunt thoracic trauma. *Ann Thorac Surg* 1967;4(2):166–170
- 6 Fu JJ, Chen CL, Wu JY. Lung torsion: survival of a patient whose hemorrhagic infarcted lung remained in situ after detorsion. *J Thorac Cardiovasc Surg* 1990;99(6):1112–1114
- 7 Daughtry DC. Traumatic torsion of the lung. *N Engl J Med* 1957;256(9):385–388
- 8 Cable DG, Deschamps C, Allen MS, et al. Lobar torsion after pulmonary resection: presentation and outcome. *J Thorac Cardiovasc Surg* 2001;122(6):1091–1093
- 9 Felson B. Lung torsion: radiographic findings in nine cases. *Radiology* 1987;162(3):631–638
- 10 Banki F, Velmahos GC. Partial pulmonary torsion after thoracotomy without pulmonary resection. *J Trauma* 2005;59(2):478–481
- 11 Parambil JG, Savci CD, Tazelaar HD, Ryu JH. Causes and presenting features of pulmonary infarctions in 43 cases identified by surgical lung biopsy. *Chest* 2005;127(4):1178–1183
- 12 Hislop A, Reid L. New pathological findings in emphysema of childhood. 2. Overinflation of a normal lobe. *Thorax* 1971;26(2):190–194
- 13 Grazia TJ, Hodges TN, Cleveland JC Jr, Sheridan BC, Zamora MR. Lobar torsion complicating bilateral lung transplantation. *J Heart Lung Transplant* 2003;22(1):102–106
- 14 Yang EM, Song ES, In Jang H, Jeong IS, Choi YY. Lung torsion after tracheoesophageal fistula repair in an infant. *Korean J Pediatr* 2013;56(4):186–190